

# BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email [editorial.bmjopen@bmj.com](mailto:editorial.bmjopen@bmj.com)

# BMJ Open

## Maintenance therapy with proton pump inhibitors and risk of gastric cancer: a nationwide population-based cohort study in Sweden

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017739
Article Type:	Research
Date Submitted by the Author:	15-May-2017
Complete List of Authors:	Brusselaers, Nele; Karolinska Institutet, Molecular medicine and Surgery Wahlin, Karl; Karolinska Institutet, Department of Molecular medicine and Surgery Engstrand, Lars; Karolinska Institutet, Microbiology, Tumour and Cell Biology Lagergren, Jesper; Karolinska Institutet, Department of Molecular medicine and Surgery; King's College London, Division of Cancer Studies
<b>Primary Subject Heading</b>:	Oncology
Secondary Subject Heading:	Epidemiology, Gastroenterology and hepatology, Oncology
Keywords:	proton pump inhibitors, chemoprevention, gastric cancer, PPI

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Title: Maintenance therapy with proton pump inhibitors and risk of gastric cancer: a nationwide population-based cohort study in Sweden**

**Running title: Proton pump inhibitors and risk of gastric cancer**

**Authors:** Nele BRUSSELAERS<sup>1</sup>, MD, PhD, Karl Wahlin<sup>1</sup>, MD, PhD, Lars ENGSTRAND<sup>2</sup>, PhD, Jesper LAGERGREN<sup>1,3</sup>, MD, PhD.

**Affiliations:** <sup>1</sup> Upper Gastrointestinal Surgery, Department of Molecular medicine and Surgery, Karolinska Institutet, Karolinska University Hospital, Stockholm, Sweden.

<sup>2</sup> Department of Microbiology, Tumor and Cell Biology and Science for Life Laboratory, Karolinska Institutet, Stockholm, Sweden.

<sup>3</sup> Division of Cancer Studies, King's College London, United Kingdom.

**Corresponding author:** Associate Professor Nele Brusselaers. Address: Department of Microbiology, Tumor and Cell Biology, Centre for Translational Microbiome Research, Karolinska Institutet, Nobels väg 16, 171 77 Stockholm, Sweden. Telephone: +46 (0)8-524 853 08. Fax: +46 (0)8-517 76280. E-mail: [nele.brusselaers@ki.se](mailto:nele.brusselaers@ki.se).

**Author contributions:** Study concept and design: All authors (NB,KW,LE, JL); Acquisition of data: NB and JL; Data collection and preparation for analyses: NB and JL; Analysis and interpretation of data: All authors (NB,KW,LE, JL); Drafting of the manuscript: NB; Critical revision of the manuscript for important intellectual content: All authors (NB,KW,LE, JL); Statistical analysis: NB and KW; Obtained funding: JL and NB.

**Word count:** 3482

**Abstract (299 words)**

*Objective:* Proton pump inhibitors (PPIs) are among the most commonly prescribed drugs. Concerns have been raised about a potentially increased risk of gastric cancer following long-term use. Our aim is to assess the risk of gastric cancer associated with PPI use, taking into account underlying indications.

*Design:* Population-based cohort study. Standardized incidence ratios (SIRs) and 95% confidence intervals (CIs) were calculated to compare the risk of gastric cancer among long-term PPI-users with the corresponding background population, while taking confounding by indication into account.

*Setting:* Population-based study in Sweden (2005-2012)

*Participants:* this study included virtually all adults residing in Sweden exposed to maintenance therapy with PPIs.

*Exposure/Intervention:* maintenance use of PPIs, defined as at least 180 days during the study period. Maintenance use of H<sub>2</sub>-receptor antagonist was evaluated for comparison reasons.

*Outcome measures:* gastric cancer (cardia and non-cardia), and subgroup analysis for gastric adenocarcinoma, as defined by the Swedish Cancer Register.

*Results:* Among 797,067 individuals on maintenance PPI therapy, the SIR of gastric cancer was over 3-fold increased (SIR=3.38, 95% CI 3.23-3.53). Increased SIRs were found in both sexes and all age groups, but were especially increased among PPI-users younger than 40 years (SIR=22.76, 95% CI 15.94-31.52). Increased SIRs were found for each indication studied,

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

including those without an association with gastric cancer, e.g. gastro-  
esophageal reflux (SIR=3.04, 95% CI 2.80-3.31), and those with a supposedly  
decreased risk, e.g. aspirin users (SIR=1.93, 95% CI 1.70-2.18). The  
association was similar for cardia and non-cardia gastric cancer. Analyses  
restricted to adenocarcinoma showed similar results to those for all gastric  
cancers. Long-term users of histamine-2 receptor antagonists, which have the  
same indications as PPIs, were not at any increased risk.

*Conclusions:* Long-term PPI-use might be an independent risk factor for  
gastric cancer. This challenges broad maintenance PPI therapy, particularly if  
the indication is weak.

## Article summary Strengths and limitations of this study

- Population-based and nationwide design based on contemporary use of proton-pump inhibitors (PPIs) – resulting in sufficient power to assess underlying indications (to assess confounding by indication).
- To our knowledge, this is the largest study to date assessing the association between PPIs and gastric cancer
- The findings are standardized for age, sex – which are often described the major confounding factors in epidemiologic studies - and calendar time. Yet, other confounders could not be taken into account because the information was not available for the background population.
- Exposure information is based on the Swedish Prescribed Drug Registry, which is initiated in July 2005 and has a complete nationwide coverage.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Introduction**

Proton pump inhibitors (PPIs), introduced in the 1980s, are potent gastric acid suppressors that reduce gastric acidity.(1-3) PPIs are among the most prescribed drugs globally,(4) used for healing peptic ulcers, counteracting gastro-esophageal reflux, eradicating *Helicobacter pylori* (in combination with antibiotics) and preventing primary or recurrent peptic ulcers, e.g. in individuals exposed to aspirin or other non-steroidal anti-inflammatory drugs (NSAIDs) or with Zollinger-Ellison syndrome (a gastrin-secreting pancreatic tumor). However, it has been suggested that long-term PPI-use increases the risk of pre-malignant gastric lesions (e.g. polyps, atrophy, and metaplasia) and gastric cancer.(2, 5, 6) Gastric acid secretion blockage may disrupt the gastric microbiome, interfere with nitrosamine formation, cause chronic atrophic gastritis and increase gastrin serum levels, which can all contribute to gastric cancer development.(2, 5, 7, 8). The effect of PPI use on the gut microbiome may even be more prominent than the effects of antibiotics.(9) Among three recent meta-analyses on the topic, one found no association between long-term PPI-use and pre-malignant gastric lesions, based on six randomized controlled trials (1789 patients in total).(2) The second included an additional trial (2343 patients in total), and found no evidence of gastric tumor development in PPI-users with atrophy or intestinal metaplasia, while an increased risk of gastric hyperplasia was indicated.(6) The third, based on 11 observational studies (94,558 participants), reported a 40% increase of gastric cancer among PPI-users.(5) However, the impact of confounding by indication remains unknown. The present study aimed to assess the risk of gastric cancer in long-term PPI-users in a population-based design, while

1  
2  
3 taking confounding by indication for such treatment into account. For  
4  
5 comparison reasons, use of histamine 2 receptor antagonists (H2RAs), which  
6  
7 are used for similar indications as PPIs, was also studied.  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Results**

***Study participants***

In total, 797,067 individuals on maintenance PPI therapy were included in the exposed cohort, resulting in 3,866,836 person-years of follow-up (mean 4.9 years). An additional 20,210 individuals had maintenance H2RA therapy, and 25,726 had both PPI and H2RA maintenance therapy. Figure 1 presents how the study participants were selected, and Table 1 shows their characteristics. Women constituted 58.5% of the PPI cohort, and a majority (66.1%) was younger than 70 years. Maintenance therapy with aspirin (34.8%) and NSAIDs (30.4%) were the most common indications for maintenance PPI use, followed by gastro-esophageal reflux (25.3%), gastro-duodenitis (13.2%) and peptic ulcer (10.0%). *Helicobacter pylori* infection was the indication in 7.3% of all participants, and dyspepsia in 5.5%. Barrett’s esophagus and Zollinger-Ellison syndrome were rare indications (<1%) and therefore not eligible for separate analysis. A third (33.4%) of the participants had more than one of the listed indications, and was therefore included in more than one indication group. Overall mortality was slightly lower among PPI users (17.3%) compared to H2RA maintenance users (19.4%), while mortality in gastric cancer patients did not differ significantly.

***Proton pump inhibitor use and overall risk of gastric cancer***

Among all participants exposed to maintenance PPIs, 2,219 (0.28%) developed gastric cancer during follow-up. Of these, 1,652 (74.4%) had non-cardia gastric cancer and 1,943 (87.6%) had adenocarcinoma (Table 1). Long-term PPI-users were at more than a 3-fold increased SIR of gastric

cancer of any type (SIR=3.38, 95% CI 3.25-3.53) and gastric adenocarcinoma (SIR=3.38, 95% CI 3.23-3.53) (Table 2). After excluding early gastric cancers, the risk remained increased (SIR=1.61, 95% CI 1.51-1.71). The risk of gastric cancer was highest among individuals receiving PPI for shorter than 1 year (SIR=12.82, 95% CI 12.19-13.47), but there was also evidence for an increased risk up to 3 years of PPI use, yet reduced risks for use over 5 years (Table 3). The strength of association was similar for cardia (SIR=3.55, 95% CI 3.27-3.86) and non-cardia gastric cancer (SIR=3.33, 95% CI 3.17-3.50). The risk of gastric cancer was similar in men (SIR=3.65, 95% CI 3.45-3.85) and women (SIR=3.07, 95% CI 2.87-3.28). The SIR was higher in younger ages; PPI-users younger than 40 years had SIR of 22.76 (95% CI 15.94-31.52) while those who were at least 70 years old had SIR of 2.76 (95% CI 2.61-2.92) (Table 2). The subgroup analyses for sex and age showed similar results as the overall findings in separate analyses for gastric adenocarcinoma, as well as for cardia and non-cardia gastric cancer (Table 2).

### ***Proton pump inhibitor use and risk of gastric cancer stratified by indication***

The SIR of gastric cancer was increased in each of the 10 studied groups of indications for PPI therapy (Table 4). The highest SIRs were found among participants exposed to indications with a known association with gastric cancer, i.e. *Helicobacter pylori* infection and peptic ulcer. However, the SIRs were also increased for indications without any such association. The SIR for gastric cancer was 3.04 (95% CI 2.80-3.31) for gastro-esophageal reflux. The

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

SIR were increased also for indications where a decreased risk of gastric cancer was expected. The SIR was 1.93 (95% CI 1.70-2.18) among those using PPIs because of aspirin use without any other indication. The SIRs were generally similar for non-cardia gastric cancer. Regarding cardia cancer, higher SIRs than for gastric cancer were found for the indication gastro-esophageal reflux and among those exposed to aspirin or other NSAIDs, while the associations for *Helicobacter pylori* infection and peptic ulcer were less strong (Table 4).

**Attributable risk**

Of all 5,823 gastric cancer patients diagnosed during the study period, 38.1% occurred among maintenance PPI-users. Based on the overall SIR of 3.38, and a prevalence of PPI-use of 10.7% among adults, the attributable fraction among PPI-users (AF) was 70.4%, and 20.3% in the total population (PAF), assuming causality. Using the lowest SIR according to Table 4 (SIR=1.41), the corresponding proportions were 29.1% (AF) and 4.2% (PAF), respectively. Among individuals younger than 40, the prevalence of PPI-use was 3.3%, and 37.1% of all 97 gastric cancers occurred among PPI-users, corresponding to AF of 95.6% and PAF of 41.8%.

**Use of histamine-2 receptor blockers only or proton pump inhibitors and histamine-2 receptor blockers and risk of gastric cancer**

There were 12 and 62 cases of gastric cancer among maintenance H2RAs users only and both H2RAs and PPIs, respectively. The risk of gastric cancer was not increased in the H2RA only group (SIR=0.57, 95% CI 0.29-0.99), and

was moderately increased in the group exposed to both PPIs and H2RAs  
(SIR=2.09, 95% CI 1.61-2.69).

For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Discussion**

This study provides some evidence of an increased risk of gastric cancer among maintenance PPI-users, including those who had PPI therapy for indications without any positive association with gastric cancer. Increased risks were found for both cardia and non-cardia gastric cancer, in both sexes and all age groups, although the risk was more pronounced in younger participants.

Strengths of the study include the nationwide and population-based design, the valid data on exposures, outcomes, and indications for PPI-use, and the large number of individuals exposed to maintenance treatment with PPIs. To our knowledge, this is the largest study to date on this topic,(10) and the cohort allowed analyses of differences in associations between underlying indications.

The main problems of this study are confounding, especially by indication, and reverse causality. Confounding by indication has been investigated by looking at the different indication groups, including groups without no increased risk of gastric cancer. Unfortunately, we could not identify any underlying indication for 25% of the PPI-users, and 39% of the H2RA-users. It is likely that clear indications were more readily recorded, while use for less rigorously diagnosed indications were more prone to be missing. This may indicate that PPI-users had more severe symptoms than H2RA-users, as also indicated by the higher proportion of PPI users presenting with multiple indications (34% vs. 19%). Yet, the lack of information on indication should not explain the associations for known indications – and those groups with no

1  
2  
3 apparent risk factors should be less likely to have severe gastrointestinal  
4  
5 symptoms. Therefore, it was unexpected to still see such increased risks of  
6  
7 gastric cancer in those groups. Unfortunately, we lacked information about  
8  
9 some potential confounders, e.g. dietary factors, obesity, tobacco smoking  
10  
11 and alcohol overconsumption since these are not collected in the nationwide  
12  
13 Health Registries. However, the lack of association between H2RAs and  
14  
15 gastric cancer argues against bias from confounding or selection from lifestyle  
16  
17 factors or any other unknown factors. Since H2RAs were used for similar  
18  
19 indications as PPIs (yet clearly became less popular), the lack of association  
20  
21 between H2RAs and gastric cancer supports that the association between  
22  
23 PPI-use and gastric cancer may be linked to the PPI medication *per se*. Yet,  
24  
25 PPIs clearly became the first choice treatment for most indications for gastric  
26  
27 acid suppression with almost 40 times more PPI maintenance users than  
28  
29 H2RA maintenance users in Sweden. The high prevalence of PPI  
30  
31 maintenance use among adults (10.7% in Sweden) also means that the large  
32  
33 majority with recognized risk factors for gastric cancer will have received PPI  
34  
35 treatment at some point. This clearly hampers assessment of PPI use as an  
36  
37 independent risk factor and this may even be a larger problem in other study  
38  
39 designs especially if indications of use cannot be assessed.  
40  
41  
42  
43  
44  
45 The problem of reverse causality, i.e. individuals taking PPIs because of  
46  
47 symptoms arising from an undetected cancer, should have been reduced by  
48  
49 only including individuals with 180 days (6 month) cumulative exposure before  
50  
51 any cancer diagnosis. In addition, we excluded all individuals who had gastric  
52  
53 cancer within a year after inclusion and also stratified the analyses by duration  
54  
55 of use; which still showed increased risks (respectively SIR=1.61 and  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

SIR=2.19 among those with an exposure duration between 1 and 3 years). These increased risks may not be entirely explained by reverse causality and detection bias alone – since most cases of gastric cancer are believed to be detected within 1 year after onset of symptoms. There are no significant waiting times or socio-economic differences in access to endoscopy in Sweden, and no obvious differences were found between tumor stages (potential detection bias),(11) or anatomical locations between the PPI-exposed cohort and the background population (Appendix 3).

The fact that the risk seems to go down with a longer-duration of use is probably because PPI is beneficial for most individuals with known risk factors for cancer – yet further research seems needed for those on maintenance therapy without gastrointestinal indications.

Although our follow-up should ideally have been longer, our maximal follow-up time is 7.5 years, which is remarkably longer than all studies included in the 2 meta-analyses assessing pre-malignant lesions based on randomized controlled trials (6-36 months as maximal follow-up).(2, 6) Only 2 smaller cohort studies had longer follow-up,(1, 10) and these were included in the only meta-analysis evaluating gastric cancer risk based on observational studies.(5) We also limited our study to maintenance use, defined as at least 180 days of exposure, which is a commonly used approach.(6) Compliance and over-the-counter availability of some PPIs (Esomeprazole, Lansoprazole and Pantoprazole – only in pharmacy, not in retail sale) may have resulted in different actual dosages. H2RA were not available over-the-counter during the study period. Therefore, and because of lack of exposure data before the

study period, we cannot be certain that the registered dosage reflects the actual dosage.

The increased risks of gastric cancer among individuals with *Helicobacter pylori* infection and peptic ulcers were anticipated, since both these conditions are risk factors for gastric cancer, but the associations were stronger than expected, suggesting an additional etiological role of the PPI-use. Importantly, there was evidence of a substantially increased risk in individuals exposed to conditions not known to increase the risk of gastric cancer, including gastro-esophageal reflux. Moreover, increased risks were found even among participants without any gastrointestinal indications who were exposed only to maintenance treatment with aspirin or other NSAIDs, although these drugs are expected to decrease the risk of gastric cancer.(12, 13) The higher risk of gastric cancer in the younger age group may be related to a recently described increase in atrophic gastritis in Sweden, as a potential consequence of the stabilizing sero-prevalence of *Helicobacter pylori* and increased prevalence of overweight and obesity.(14)

The difference in the association with gastric cancer following maintenance therapy with PPIs and H2RAs could be explained simply by the fact that PPIs are more potent than H2RAs in inhibiting acid secretion.(15) PPIs block the gastric proton pumps, while H2RAs compete with histamine for the histamine-2 receptors. At recommended dosages, PPIs induce a more profound and prolonged acid inhibition than H2RAs. Moreover, the acid inhibition of PPIs also increases over time, while the effects of H2RAs fade.(15)



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

There are several mechanisms that might explain an association between PPI-use and gastric cancer. Histopathological changes occur due to the blockage of the normal gastric acid secretion, leading to over-stimulation (hypergastrinemia) that might cause hyperproliferation of the gastric mucosa, chronic hypochlorhydria (reduction of hydrochloric acid in the gastric juice), chronic inflammation and disappearance of normal mucosal glands and their replacement by intestinal glands, and possible gastric atrophy.(6) All these mucosal changes could promote gastric carcinogenesis. The blocked gastric acid secretion could decrease physiological defense mechanisms against pathogenic bacteria.(7, 8) Diarrhea is for example a well-known side-effect of PPI-use, which is often due to infection with the bacteria *Clostridium difficile*, non-thyphoid *Salmonella species* or *Campylobacter jejuni*.(7) This reduced bacterial defense mechanism might, in the long term, result in chronic inflammation and eventually gastric cancer development. Decreased gastric acidity may also result in increased bacterial colonization, including non-gastric microorganisms, and a greater number of bacteria that produce nitrosamines, which are well-established carcinogens of the gastric mucosa.(5, 16)

The results of a single study, even of this size, cannot determine causality. Yet, the effect size, consistency across indications (especially those not associated with an increased risk of gastric cancer), lack of association with H2RAs, and plausible carcinogenic mechanisms mean that maintenance PPI-use cannot be dismissed as a potentially independent risk factor for gastric

1  
2  
3 cancer. A higher awareness for gastric cancer could potentially be considered  
4  
5 in maintenance users of PPI, especially if known underlying risk factors are  
6  
7 present. Thus, these findings need confirmation in future investigations,  
8  
9 especially considering the generalizability of the results to populations with  
10  
11 higher incidence of gastric cancer or other distribution of risk factors.  
12  
13

14  
15  
16 To conclude, this large and population-based cohort study provides evidence  
17  
18 of a substantially increased risk of gastric cancer following maintenance use  
19  
20 of PPIs *per se*. The consistency across sexes, age groups, and indication  
21  
22 groups, including indications that do not increase the risk of gastric cancer,  
23  
24 support the overall finding. If confirmed in further research, especially among  
25  
26 those without gastrointestinal risk factors, these findings challenge a broad  
27  
28 use of long-term PPI-use, particularly for conditions where the indication is  
29  
30 weak.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Methods**

***Design***

This was a nationwide Swedish population-based cohort study designed to examine the risk of gastric cancer in individuals exposed to maintenance therapy with PPIs (and to maintenance use of H2RAs), compared to the Swedish background population of the same sex, age and calendar period (7.1-7.6 million adults).(17)

Only adults (at least 18 years) without a history of any cancer were included. The participants were followed up from the first prescription of a PPI (or H2RA) during the period 1<sup>st</sup> July 2005 to 31<sup>st</sup> December 2012. The data were derived from high-quality and nationwide Swedish registries and information on individuals was linked between the registries by means of the unique Swedish personal identity number.(18) The study cohort included all Swedish residents who received at least one dispensed prescription of commonly prescribed drugs (listed in Appendix 1) between 1<sup>st</sup> July 2005 and 31<sup>st</sup> December 2014 (with follow-up for cancer until 31<sup>st</sup> December 2012). The study was approved by the Regional Ethical Review Board in Stockholm, and informed consent was not required (2014/1291-31/4).

***Patient involvement***

The Swedish patient organization for cancer of the esophagus, stomach, liver, and pancreas was involved in supporting the present study ([www.palema.org](http://www.palema.org)). The development of the research question and outcome measures were informed by patients' priorities, experience, and preferences. The results will be disseminated to study participants by means of patient

organizations. Patients are thanked in the acknowledgements.

### **Exposure**

The study exposure was maintenance therapy with a PPI (or an H2RA) according to the Swedish Prescribed Drug Registry, defined as a cumulative defined daily dose (DDD) of at least 6 months ( $\geq 180$  days) during the study period (before a potential cancer diagnosis). The DDD was the average maintenance dose per day for a drug used for its main indication in adults, which follows the World Health Organization (WHO) definition. This cumulative DDD was estimated by adding the DDD per package, which takes both the potency and the quantity of the drug into account, and is therefore a proxy for the duration of the exposure. The Anatomical Therapeutic Chemical classification system (ATC) was used to identify codes representing PPIs (code A02BC) and H2RAs (code A02BA). The participants were divided into 3 mutually exclusive exposure groups: (1) PPI-users ( $\geq 180$  days on PPIs and  $< 180$  days on H2RAs); (2) H2RA users ( $\geq 180$  days on H2RAs and  $< 180$  days on PPIs); and (3) PPI and H2RA users ( $\geq 180$  days on PPIs and  $\geq 180$  days on H2RAs). These medications were also available over-the-counter in Sweden, but only in small packages and at a much higher price per dose.<sup>(19)</sup> Thus, maintenance users had their medication prescribed and were therefore included in this study.

### **Outcome**

The main outcome was a first episode of any gastric cancer according to the Swedish Cancer Registry. Gastric cancer was defined by the diagnosis codes

C16.0 (cardia) and C16.1-C16.9 (non-cardia) in the 10<sup>th</sup> version of the International Classification of Diseases (ICD), and adenocarcinoma (dominating histological type) was defined by the histology code 096.

**Confounders**

Age, sex, and calendar period were adjusted for in the design. Confounding by indication was evaluated by analyzing indications for PPI-use separately (see Appendix 2 for ICD and ATC codes): gastro-esophageal reflux disease, Barrett’s esophagus, peptic ulcer disease, Zollinger-Ellison syndrome, gastro-duodenitis, dyspepsia/disruption of gastric function, *Helicobacter pylori* infection, *Helicobacter pylori* eradication (combined with *Helicobacter pylori* infection in the analyses), and maintenance use (≥180 days) of aspirin or other NSAIDs.

**Data sources**

Four nationwide Swedish registries, all maintained by the governmental National Board of Health and Welfare, provided data for the study.

**The Swedish Prescribed Drug Registry** started in 1<sup>st</sup> July 2005, and includes all drugs prescribed and dispensed in Sweden. Information from these prescriptions is transferred for registration monthly.(20) This registry is highly complete (patient identity data are missing in <0.3% of all items),(20) and was used to collect data on type of medication, dates of prescribing and dispensing, and DDD per package.

**The Swedish Cancer Registry** has at least a 96% complete registration of all cancers in Sweden since1958,(21) with 98% completeness for gastric

1  
2  
3 cancer.(22, 23) This registry was used to identify gastric cancer cases among  
4 the cohort members and the corresponding background population (exact  
5 annual numbers were available for the used age and sex categories), and to  
6 exclude individuals with a history of any cancer.  
7  
8  
9

10  
11 **The Swedish Patient Registry** has a full nationwide coverage for all  
12 diagnoses from 1987.(18) The registration of out-patient specialist care is  
13 nationwide since 2001. This registry was used to collect information on  
14 indications for PPI-use.  
15  
16  
17  
18  
19

20  
21 **The Swedish Causes of Death Registry** has 100% completeness for  
22 recording of dates of death since 1952.(24) This registry was used to collect  
23 date of death.  
24  
25  
26  
27  
28  
29

### 30 **Statistical analyses**

31 All analyses were conducted according to a pre-planned protocol. The risk of  
32 gastric cancer was compared between the exposed cohort and the entire  
33 Swedish background population of the same sex (male or female), age group  
34 (categorized as 18-39, 40-49, 50-59, 60-69 or ≥70 years), and calendar period  
35 (2005-2006, 2007-2009 or 2010-2012). Standardized incidence ratios (SIRs)  
36 and 95% confidence intervals (CIs) were calculated by dividing the observed  
37 number of gastric cancer cases with the expected number, accounting for  
38 changes in age and calendar categories.(25) The expected numbers were  
39 derived from the Swedish Cancer Registry and Statistics Sweden.(17) Follow-  
40 up time was calculated from the dispense date of the first prescription of PPI  
41 (or H2RA) within the study period, until death, any cancer, or end of the study  
42 period (December 31, 2012), whichever occurred first. Subgroup analyses  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

were stratified for sex and age. Stratified analyses were performed for each indication whenever at least 10,000 exposed individuals were identified with this indication. To assess reverse causality (protopathic bias), a sensitivity analysis was conducted excluding all cancer cases occurring within one year of the start of the study. Duration of PPI-use was estimated based on the total DDD per package prescribed before any cancer diagnosis, and was categorized into <1.0 year, 1.0-2.9 years, 3.0-4.9 years or ≥5 years. The attributable fraction (AF) and the population attributable fraction (PAF) were calculated to estimate the proportion of gastric cancer in PPI-users and the total population, respectively (assuming causality). The following formulas were used:  $AF = (SIR - 1) / SIR$  and  $PAF = p(SIR - 1) / (p(SIR - 1) + 1)$ , where p represents the prevalence of PPI maintenance use in the population (based on the population size in 2009).<sup>(17)</sup> There were no missing data on exposures, outcomes, age, sex or calendar period. When no information was available for the indication for PPI-use (25.0%), the indication was considered absent.

**Acknowledgements:** We are grateful to all thousands of patients, physicians and healthcare staff members who contributed to the data collection, and to the Swedish National Board of Health and Welfare that collected the data. NB had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. NB conducted and is responsible for the data-analysis. Everyone who contributed significantly to the work is listed as co-author.

**Funding source:** The work was supported by the Swedish Research Council (839-2008-7496), Swedish Cancer Society (CAN 2015/460), the Karolinska Institutet Distinguished Professor Award to Jesper Lagergren (D-02418/2010) and SFO Epidemiology (Young Scholar Grant to Nele Brusselaers).

**Competing interest statement:** There are no competing interests.

**Data sharing statement:** We are willing to share data upon request after ethical approval has been approved by the relevant committee and the governmental agencies that maintain the data.

## References

1. Moller H, Nissen A, Mosbech J. Use of cimetidine and other peptic ulcer drugs in Denmark 1977-1990 with analysis of the risk of gastric cancer among cimetidine users. *Gut*. 1992 Sep;33(9):1166-9. PubMed PMID: 1358764. Pubmed Central PMCID: PMC1379479.
2. Eslami L, Nasser-Moghaddam S. Meta-analyses: does long-term PPI use increase the risk of gastric premalignant lesions? *Arch Iran Med*. 2013 Aug;16(8):449-58. PubMed PMID: 23906249. Epub 2013/08/03. eng.
3. Shin JM, Sachs G. Pharmacology of proton pump inhibitors. *Curr Gastroenterol Rep*. 2008 Dec;10(6):528-34. PubMed PMID: 19006606. Pubmed Central PMCID: PMC2855237.
4. The World Medicines Situation (WMS) 2011 - Pharmaceutical consumption - Annex 4 Top 5 medicine classes by volume and country income category.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

5. Ahn JS, Eom CS, Jeon CY, Park SM. Acid suppressive drugs and gastric cancer: a meta-analysis of observational studies. *World J Gastroenterol*. 2013 Apr 28;19(16):2560-8. PubMed PMID: 23674860. Pubmed Central PMCID: 3646149. Epub 2013/05/16. eng.

6. Song H, Zhu J, Lu D. Long-term proton pump inhibitor (PPI) use and the development of gastric pre-malignant lesions. *Cochrane Database Syst Rev*. 2014;12:CD010623. PubMed PMID: 25464111.

7. Canani RB, Terrin G. Gastric acidity inhibitors and the risk of intestinal infections. *Curr Opin Gastroenterol*. 2010 Jan;26(1):31-5. PubMed PMID: 19907324.

8. Seto CT, Jeraldo P, Orenstein R, Chia N, DiBaise JK. Prolonged use of a proton pump inhibitor reduces microbial diversity: implications for *Clostridium difficile* susceptibility. *Microbiome*. 2014;2:42. PubMed PMID: 25426290. Pubmed Central PMCID: PMC4242847.

9. Imhann F, Bonder MJ, Vich Vila A, Fu J, Mujagic Z, Vork L, et al. Proton pump inhibitors affect the gut microbiome. *Gut*. 2016 May;65(5):740-8. PubMed PMID: 26657899. Pubmed Central PMCID: 4853569.

10. Poulsen AH, Christensen S, McLaughlin JK, Thomsen RW, Sorensen HT, Olsen JH, et al. Proton pump inhibitors and risk of gastric cancer: a population-based cohort study. *Br J Cancer*. 2009 May 5;100(9):1503-7. PubMed PMID: 19352380. Pubmed Central PMCID: PMC2694435.

11. Sobin L, Wittekind C. International Union Against Cancer (UICC)- 6th edition: Classification of malignant tumours. 6th ed2002.

12. Gurpinar E, Grizzle WE, Piazza GA. NSAIDs inhibit tumorigenesis, but how? Clin Cancer Res. 2014 Mar 1;20(5):1104-13. PubMed PMID: 24311630. Pubmed Central PMCID: PMC3947450.
13. Ford AC. Chemoprevention for gastric cancer. Best Pract Res Clin Gastroenterol. 2011 Aug;25(4-5):581-92. PubMed PMID: 22122773. Epub 2011/11/30. eng.
14. Song H, Held M, Sandin S, Rautelin H, Eliasson M, Soderberg S, et al. Increase in the Prevalence of Atrophic Gastritis Among Adults Age 35 to 44 Years Old in Northern Sweden Between 1990 and 2009. Clin Gastroenterol Hepatol. 2015 Sep;13(9):1592-600 e1. PubMed PMID: 25857683.
15. Lamers CB. The changing role of H2-receptor antagonists in acid-related diseases. Eur J Gastroenterol Hepatol. 1996 Oct;8 Suppl 1:S3-7. PubMed PMID: 8930573.
16. Engstrand L, Lindberg M. Helicobacter pylori and the gastric microbiota. Best Pract Res Clin Gastroenterol. 2013 Feb;27(1):39-45. PubMed PMID: 23768551. Epub 2013/06/19. eng.
17. Statistics Sweden - Statistical Database Population Statistics (accessed 06 April 2016) <http://www.statistikdatabasen.scb.se/>. Available from: <http://www.statistikdatabasen.scb.se/>.
18. Ludvigsson JF, Andersson E, Ekbom A, Feychting M, Kim JL, Reuterwall C, et al. External review and validation of the Swedish national inpatient register. BMC Public Health. 2011;11:450. PubMed PMID: 21658213. Pubmed Central PMCID: 3142234. Epub 2011/06/11. eng.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

19. FASS (Farmaceutiska specialiteter i Sverige/ Pharmaceutical specialities in Sweden) [Internet]. 2015 [cited November 2016]. Available from: [www.fass.se](http://www.fass.se).

20. Wettermark B, Hammar N, Fored CM, Leimanis A, Otterblad Olausson P, Bergman U, et al. The new Swedish Prescribed Drug Register--opportunities for pharmacoepidemiological research and experience from the first six months. *Pharmacoepidemiol Drug Saf*. 2007 Jul;16(7):726-35. PubMed PMID: 16897791. Epub 2006/08/10. eng.

21. Barlow L, Westergren K, Holmberg L, Talback M. The completeness of the Swedish Cancer Register: a sample survey for year 1998. *Acta Oncol*. 2009;48(1):27-33. PubMed PMID: 18767000. Epub 2008/09/04. eng.


22. Lindblad M, Ye W, Lindgren A, Lagergren J. Disparities in the classification of esophageal and cardia adenocarcinomas and their influence on reported incidence rates. *Ann Surg*. 2006 Apr;243(4):479-85. PubMed PMID: 16552198. Pubmed Central PMCID: 1448962. Epub 2006/03/23. eng.

23. Ekstrom AM, Signorello LB, Hansson LE, Bergstrom R, Lindgren A, Nyren O. Evaluating gastric cancer misclassification: a potential explanation for the rise in cardia cancer incidence. *J Natl Cancer Inst*. 1999 May 5;91(9):786-90. PubMed PMID: 10328109. Epub 1999/05/18. eng.

24. The National Board of Health and Welfare, Sweden. Causes of death in Sweden 2009. 2011. Available from: <http://www.socialstyrelsen.se/publikationer2011/2011-3-22>.

25. Breslow N, Day N. Statistical Methods in Cancer Research: The design and analysis of cohort studies. Lyon1987.

**Figure 1: Flowchart of the construction of the cohort exposed to maintenance therapy with proton pump inhibitors (PPIs) and/or H<sub>2</sub> receptor antagonists (H<sub>2</sub>RAs).**



**Table 1: Descriptive characteristics of the study cohort exposed to maintenance therapy with proton pump inhibitors (PPIs) and/or H2 receptor antagonists (H2RAs).**

		PPIs only Number (%)	H2RAs only Number (%)	PPIs and H2RAs Number (%)
<b>Total</b>		797,067	20,210	25,726
<b>Sex</b>				
	<i>Men</i>	330,885 (41.5)	8,076 (40.0)	9,778 (38.0)
	<i>Women</i>	466,182 (58.5)	12,134 (60.0)	15,948 (62.0)
<b>Age (years)</b>				
	<40	89,231 (11.2)	1,885 (9.3)	1,822 (7.1)
	40-49	104,003 (13.1)	2,461 (12.2)	3,268 (12.7)
	50-59	155,963 (19.6)	3,971 (19.7)	5,878 (22.9)
	60-69	177,606 (22.3)	4,563 (22.6)	6,354 (24.7)
	≥70	270,264 (33.9)	7,330 (36.3)	8,404 (32.7)
<b>Calendar period</b>				
	2005-2006	446,068 (56.0)	16,550 (81.9)	23,066 (90.0)
	2007-2009	222,257 (27.9)	2,670 (13.2)	2,304 (9.0)
	2010-2012	128,742 (16.2)	990 (4.9)	356 (1.4)
<b>Indications</b>				
	<i>Gastro-esophageal reflux</i>	201,868 (25.3)	2,073 (10.3)	6,727 (26.2)
	<i>Barrett's esophagus</i>	6,044 (0.8)	17 (0.1)	172 (0.7)
	<i>Peptic ulcers</i>	79,597 (10.0)	1,226 (6.1)	2,633 (10.2)
	<i>Zollinger-Ellison syndrome</i>	31 (0.0)	0 (0.0)	3 (0.01)
	<i>Gastro-duodenitis</i>	104,955 (13.2)	1,321 (6.5)	3,695 (14.4)
	<i>Dyspepsia</i>	43,901 (5.5)	649 (3.2)	1,664 (6.5)
	<i>Helicobacter pylori</i>	58,366 (7.3)	641 (3.2)	2,041 (7.9)
	<i>Long-term NSAID use</i>	241,958 (30.4)	5,012 (24.8)	9,628 (37.4)
	<i>Long-term aspirin use</i>	277,128 (34.8)	6,533 (32.3)	9,929 (38.6)
<b>Number of indications</b>				
	0	199,608 (25.0)	7,919 (39.2)	5,402 (21.0)
	1	330,027 (41.4)	8,487 (42.0)	10,320 (40.1)
	≥2	267,432 (33.6)	3,804 (18.8)	10,004 (38.9)
<b>Gastric cancer</b>				
	<i>All</i>	2,219 (0.28)	12 (0.06)	62 (0.24)
	<i>Adenocarcinoma</i>	1,943 (0.24)	10 (0.05)	54 (0.21)
	<i>Cardia</i>	567 (0.07)	1 (0.00)	13 (0.05)
	<i>Non-cardia</i>	1,652 (0.21)	11 (0.05)	49 (0.19)
<b>Years of follow-up</b>				
	Total	3,866,836	116,015	163,519
	Mean	4.9	5.7	6.4
<b>Cumulative defined daily dosages (DDD), estimated number of days</b>				
	Median	856	450	1741
	Interquartile range	356-2102	270-990	1043-2732
<b>Mortality</b>				
	Total	137,619 (17.3)	3,915 (19.4)	3,812 (14.8)
	In gastric cancer cases	1,473 (66.4)	9 (75.0)	39 (62.9)

**Table 2: Standardized incidence ratios (SIRs) by age, sex and calendar period and 95% confidence intervals (CIs) of gastric cancer in all individuals exposed to proton pump inhibitors (PPIs), stratified by sex and age-group.**

	Gastric cancer		Gastric adenocarcinoma		Cardia cancer		Non-cardia gastric cancer	
	<i>Number of cases</i>	<i>SIR (95% CI)</i>	<i>Number of cases</i>	<i>SIR (95% CI)</i>	<i>Number of cases</i>	<i>SIR (95% CI)</i>	<i>Number of cases</i>	<i>SIR (95% CI)</i>
Total	2,219	3.38 (3.25-3.53)	1,943	3.38 (3.23-3.53)	567	3.55 (3.27-3.86)	1,652	3.33 (3.17-3.50)
Sex								
Men	1,296	3.65 (3.45-3.85)	1,166	3.66 (3.45-3.88)	437	3.84 (3.48-4.21)	859	3.56 (3.33-3.81)
Women	923	3.07 (2.87-3.28)	777	3.02 (2.82-3.25)	130	2.84 (2.38-3.38)	793	3.11 (2.99-3.34)
Age, years								
<40	36	22.76 (15.94-31.52)	27	25.30 (16.67-36.80)	7	27.33 (10.95-56.30)	29	21.88 (14.65-31.43)
40-49	94	7.36 (5.95-9.01)	76	7.76 (6.11-9.71)	24	7.54 (4.83-11.22)	70	7.30 (5.69-9.23)
50-59	303	5.63 (5.01-6.30)	263	5.82 (5.14-6.57)	94	5.50 (4.44-6.72)	209	5.69 (4.94-6.51)
60-69	576	3.85 (3.54-4.18)	501	3.81 (3.49-4.16)	191	3.94 (3.40-4.54)	385	3.81 (3.44-4.21)
≥70	1,210	2.76 (2.61-2.92)	1,076	2.77 (2.61-2.94)	251	2.77 (2.44-3.14)	959	2.76 (2.59-2.94)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Table 3: Standardized incidence ratios (SIRs) by estimated duration of use and 95% confidence intervals (CIs) of gastric cancer in all individuals exposed to proton pump inhibitors (PPIs)**

Duration	Number of cases /	
	Total	SIR (95% CI)
< 1.0 year	1,552	12.82 (12.19-13.47)
1.0-2.9 years	2,193	2.19 (1.98-2.42)
3.0-4.9 years	1,098	1.10 (0.91-1.31)
≥ 5.0 years	153	0.61 (0.52-0.72)

**Table 4: Standardized incidence ratios (SIRs) by age, sex and calendar period and 95% confidence intervals (CIs) of gastric cancer in all individuals exposed to proton pump inhibitors (PPIs), stratified by indication.**

	Gastric cancer		Cardia cancer		Non-cardia gastric cancer	
	Number of cases / total	SIR (95% CI)	Number of cases / total	SIR (95% CI)	Number of cases / total	SIR (95% CI)
All	2,219/ 797,067	3.38 (3.25-3.53)	567/ 797,067	3.55 (3.27-3.86)	1652/ 797,067	3.33 (3.17-3.50)
Gastro-esophageal reflux	557/ 201,868	3.04 (2.80-3.31)	175/ 201,868	3.84 (3.29-4.45)	382/ 201,868	2.78 (2.51-3.07)
Peptic ulcers	721/ 79,597	8.75 (8.12-9.41)	71/ 79,597	3.51 (2.74-4.43)	650/ 79,597	10.45 (9.66-11.28)
Gastro-duodenitis	350/ 104,955	3.68 (3.31-4.09)	63/ 104,955	2.71 (2.09-3.47)	287/ 104,955	3.99 (3.55-4.48)
Dyspepsia	136/ 43,901	3.07 (2.58-3.63)	33/ 43,901	3.10 (2.13-4.35)	103/ 43,901	3.07 (2.50-3.72)
<i>Helicobacter pylori</i>	440/ 58,366	9.76 (8.87-10.71)	48/ 58,366	4.19 (3.09-5.55)	392/ 58,366	11.67 (10.53-12.87)
No <i>Helicobacter pylori</i>	1,779/ 781,955	2.91 (2.78-3.05)	519/ 781,955	3.50 (3.21-3.82)	1,302/ 781,955	2.73 (2.58-2.88)
NSAIDs exposure	347/ 256,598	1.82 (1.64-2.03)	94/ 256,598	2.07 (1.67-2.53)	253/ 256,598	1.75 (1.54-1.97)
Only NSAIDs exposure	93/ 118,987	1.41 (1.14-1.73)	36/ 118,987	2.29 (1.60-3.17)	57/ 118,987	1.14 (0.86-1.48)
Aspirin exposure	872/ 293,590	2.42 (2.26-2.60)	211/ 293,590	2.73 (2.37-3.13)	588/ 293,590	2.33 (2.14-2.53)
Only aspirin exposure	252/ 126,883	1.93 (1.70-2.18)	85/ 126,883	2.73 (2.18-3.37)	169/ 126,883	1.68 (1.43-1.95)

\* NSAIDs, non-steroidal anti-inflammatory drugs. One individual could have different indications, except for the group exposed only to NSAIDs

or aspirin.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Appendix 1: Description of original cohort, the “Chemoprevention of Cancer” cohort.**

This cohort included all individuals residing in Sweden who received at least one dispensed prescription of one of the following commonly prescribed drugs between July 2005 and December 2014 (with corresponding ATC codes): sex hormones (G03), drugs for peptic ulcers and gastro-esophageal reflux disease (A02B), acetylsalicylic acid (B01AC06, N02BA), non-steroidal anti-inflammatory drugs (M01A), HMG CoA reductase inhibitors (C10AA, C10BA), drugs affecting bone structure and mineralization, (M05B), and antibiotics (J01AA, J01CA04, J01FA, J01MA, J01XD, J01XE, J04AB04). This cohort included approximately 85% of all Swedish residents, with especially high coverage of adults. All adult individuals receiving at least one dispensed prescription of a gastric acid inhibitor (proton pump inhibitor or H2 receptor antagonist) were included.

**Appendix 2: International Classification of Diseases (ICD) and Anatomical Therapeutic Chemical classification system (ATC) codes to describe the indication for gastric acid inhibitors.**

	<b>ICD-7</b> (Since 1955)	<b>ICD-8</b> (Since 1968)	<b>ICD-9</b> (Since 1987)	<b>ICD-10</b> (Since 1997)
Reflux codes	784,3; 539,11- 539,12; 560,4	530,93- 530,94; 551,3; 784,3	530B-C; 553D; 787B	K20-21; K44; R12
Barrett's esophagus	-	-	-	K227
Peptic ulcer disease	540-541	531-533	531-533	K25-K27
Zollinger-Ellison syndrome / Hypergastrinemia	-	-	251F	E16.4
Gastro-duodenitis	543	535	535	K29
Dyspepsia/disruption of gastric function	544	536	536	K30-K31
<i>Helicobacter pylori</i> infection	-	-	-	B96.8 or B98.0
<b>ATC</b>				
Non-steroidal anti-inflammatory drugs	M01A			
Aspirin	B01AC06, N02BA			
<i>Helicobacter pylori</i> eradication	A02BD, or A02BC (PPI) in combination with 2 out of 3 antibiotics (Amoxicillin [J01CA04]; Clarithromycin [J01FA09], or metronidazole [J01XD]); or combined with metronidazole [J01XD], doxycycline [J01AA02] and bismuth [A02BX05]; or with amoxicillin [J01CA04] and levofloxacin [J01MA12].			

Note: an indication was considered present if the ICD code occurred at least once in the in- or out-Patient Registry. NSAIDs and Aspirin were only considered if the cumulative dosage was  $\geq 180$  days.

Appendix 3: Tumor stage and anatomical locations of gastric cancer in the cohort exposed to maintenance use of proton pump inhibitors (PPIs) and in the Swedish background population of the same age.

	PPI maintenance cohort	Background population
	Number (%)	Number (%)
<b>Total</b>	2219 (100.0)	5821 (100.0)
<b>Tumor stage - Cardia</b>		
0-1	72 (12.7)	134 (8.9)
2	36 (6.4)	80 (5.3)
3	207 (36.5)	480 (31.9)
4	145 (25.6)	496 (33.0)
unclear/incomplete	107 (18.9)	315 (20.9)
<b>Tumor stage - Non-cardia</b>		
0-1	291 (17.6)	617 (14.3)
2	68 (4.1)	168 (3.9)
3	408 (24.7)	979 (22.7)
4	422 (25.5)	1260 (29.2)
unclear/incomplete	463 (28.0)	1292 (29.9)
<b>Anatomical location</b>		
Cardia	567 (25.6)	1505 (25.9)
Fundus	97 (4.4)	268 (4.6)
Corpus	389 (17.5)	1040 (17.9)
Antrum	398 (17.9)	915 (15.7)
Pylorus	157 (7.1)	303 (5.2)
Minor curvature	74 (3.3)	173 (3.0)
Major curvature	33 (1.5)	93 (1.6)
Overgrowth of unknown origin	2 (0.1)	15 (0.3)
Unspecified	502 (22.6)	1509 (25.9)

STROBE Statement—Checklist of items that should be included in reports of *cohort studies*

	Item No	Recommendation	Page
Title and abstract	1	(a) Indicate the study’s design with a commonly used term in the title or the abstract	p 1-2 (abstract & title)
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	p 2
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	p 4-5
Objectives	3	State specific objectives, including any prespecified hypotheses	p 4-5
Methods			
Study design	4	Present key elements of study design early in the paper	p 6-10
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	p 6-10
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	p 6-10
		(b) For matched studies, give matching criteria and number of exposed and unexposed	n/a
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	p 6-10 + appendix
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	p 6-10
Bias	9	Describe any efforts to address potential sources of bias	p 6-10
Study size	10	Explain how the study size was arrived at	Fig 1
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	p 6-10
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	p 9-10
		(b) Describe any methods used to examine subgroups and interactions	p 9-10
		(c) Explain how missing data were addressed	p 9-10
		(d) If applicable, explain how loss to follow-up was addressed	n/a
		(e) Describe any sensitivity analyses	p 9-10
Results			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Fig 1
		(b) Give reasons for non-participation at each stage	Fig 1
		(c) Consider use of a flow diagram	Fig 1
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	p 11, Table 1
		(b) Indicate number of participants with missing data for each variable of interest	n/a
		(c) Summarise follow-up time (eg, average and total amount)	Table 1
Outcome data	15*	Report numbers of outcome events or summary measures over time	Tables 1-3
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted	Tables 2-3

		estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	
		(b) Report category boundaries when continuous variables were categorized	Table 1-2
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	Discussion p 13
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	p 11-14
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	p 15-19
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	p 15-19
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	p 15-19
Generalisability	21	Discuss the generalisability (external validity) of the study results	p 15-19
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	p 3, p 10

\*Give information separately for exposed and unexposed groups.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at <http://www.strobe-statement.org>.

# BMJ Open

## Maintenance therapy with proton pump inhibitors and risk of gastric cancer: a nationwide population-based cohort study in Sweden

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017739.R1
Article Type:	Research
Date Submitted by the Author:	31-Aug-2017
Complete List of Authors:	Brusselaers, Nele; Karolinska Institutet, Molecular medicine and Surgery Wahlin, Karl; Karolinska Institutet, Department of Molecular medicine and Surgery Engstrand, Lars; Karolinska Institutet, Microbiology, Tumour and Cell Biology Lagergren, Jesper; Karolinska Institutet, Department of Molecular medicine and Surgery; King's College London, Division of Cancer Studies
<b>Primary Subject Heading</b>:	Oncology
Secondary Subject Heading:	Epidemiology, Gastroenterology and hepatology, Oncology
Keywords:	proton pump inhibitors, chemoprevention, gastric cancer, PPI

SCHOLARONE™  
Manuscripts

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Title: Maintenance therapy with proton pump inhibitors and risk of gastric cancer: a nationwide population-based cohort study in Sweden**

**Running title: Proton pump inhibitors and risk of gastric cancer**

**Authors:** Nele BRUSSELAERS<sup>1</sup>, MD, PhD, Karl Wahlin<sup>1</sup>, MD, PhD, Lars ENGSTRAND<sup>2</sup>, PhD, Jesper LAGERGREN<sup>1,3</sup>, MD, PhD.

**Affiliations:** <sup>1</sup> Upper Gastrointestinal Surgery, Department of Molecular medicine and Surgery, Karolinska Institutet, Karolinska University Hospital, Stockholm, Sweden.

<sup>2</sup> Department of Microbiology, Tumor and Cell Biology and Science for Life Laboratory, Karolinska Institutet, Stockholm, Sweden.

<sup>3</sup> Division of Cancer Studies, King's College London, United Kingdom.

**Corresponding author:** Associate Professor Nele Brusselaers. Address: Department of Microbiology, Tumor and Cell Biology, Centre for Translational Microbiome Research, Karolinska Institutet, Nobels väg 16, 171 77 Stockholm, Sweden. Telephone: +46 (0)8-524 853 08. Fax: +46 (0)8-517 76280. E-mail: [nele.brusselaers@ki.se](mailto:nele.brusselaers@ki.se).

**Author contributions:** Study concept and design: All authors (NB,KW,LE, JL); Acquisition of data: NB and JL; Data collection and preparation for analyses: NB and JL; Analysis and interpretation of data: All authors (NB,KW,LE, JL); Drafting of the manuscript: NB; Critical revision of the manuscript for important intellectual content: All authors (NB,KW,LE, JL); Statistical analysis: NB and KW; Obtained funding: JL and NB.

**Word count:** 3699

**Abstract (299 words)**

*Objective:* Proton pump inhibitors (PPIs) are among the most commonly prescribed drugs. Concerns have been raised about a potentially increased risk of gastric cancer following long-term use. Our aim is to assess the risk of gastric cancer associated with PPI use, taking into account underlying indications.

*Design:* Population-based cohort study. Standardized incidence ratios (SIRs) and 95% confidence intervals (CIs) were calculated to compare the risk of gastric cancer among long-term PPI-users with the corresponding background population, while taking confounding by indication into account.

*Setting:* Population-based study in Sweden (2005-2012)

*Participants:* this study included virtually all adults residing in Sweden exposed to maintenance therapy with PPIs.

*Exposure/Intervention:* maintenance use of PPIs, defined as at least 180 days during the study period. Maintenance use of H<sub>2</sub>-receptor antagonist was evaluated for comparison reasons.

*Outcome measures:* gastric cancer (cardia and non-cardia), and subgroup analysis for gastric adenocarcinoma, as defined by the Swedish Cancer Register.

*Results:* Among 797,067 individuals on maintenance PPI therapy, the SIR of gastric cancer was over 3-fold increased (SIR=3.38, 95% CI 3.23-3.53). Increased SIRs were found in both sexes and all age groups, but were especially increased among PPI-users younger than 40 years (SIR=22.76, 95% CI 15.94-31.52). Increased SIRs were found for each indication studied,



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

including those without an association with gastric cancer, e.g. gastro-  
esophageal reflux (SIR=3.04, 95% CI 2.80-3.31), and those with a supposedly  
decreased risk, e.g. aspirin users (SIR=1.93, 95% CI 1.70-2.18). The  
association was similar for cardia and non-cardia gastric cancer. Analyses  
restricted to adenocarcinoma showed similar results to those for all gastric  
cancers. Long-term users of histamine-2 receptor antagonists, which have the  
same indications as PPIs, were not at any increased risk.

*Conclusions:* Long-term PPI-use might be an independent risk factor for  
gastric cancer. This challenges broad maintenance PPI therapy, particularly if  
the indication is weak.

## Article summary Strengths and limitations of this study

- Population-based and nationwide design based on contemporary use of proton-pump inhibitors (PPIs) – resulting in sufficient power to assess underlying indications (to assess confounding by indication).
- To our knowledge, this is the largest study to date assessing the association between PPIs and gastric cancer
- The findings are standardized for age, sex – which are often described the major confounding factors in epidemiologic studies - and calendar time. Yet, other confounders could not be taken into account because the information was not available for the background population.
- Exposure information is based on the Swedish Prescribed Drug Registry, which is initiated in July 2005 and has a complete nationwide coverage.
- Although this study provides some evidence for an association between PPI use and gastric cancer beyond gastro-intestinal indications of use, conclusions considering causality can never be drawn based on a single study.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Introduction**

Proton pump inhibitors (PPIs), introduced in the 1980s, are potent gastric acid suppressors that reduce gastric acidity.(1-3) PPIs are among the most prescribed drugs globally,(4) used for healing peptic ulcers, counteracting gastro-esophageal reflux, eradicating *Helicobacter pylori* (in combination with antibiotics) and preventing primary or recurrent peptic ulcers, e.g. in individuals exposed to aspirin or other non-steroidal anti-inflammatory drugs (NSAIDs) or with Zollinger-Ellison syndrome (a gastrin-secreting pancreatic tumor). However, it has been suggested that long-term PPI-use increases the risk of pre-malignant gastric lesions (e.g. polyps, atrophy, and metaplasia) and gastric cancer.(2, 5, 6) Gastric acid secretion blockage may disrupt the gastric microbiome, interfere with nitrosamine formation, cause chronic atrophic gastritis and increase gastrin serum levels, which can all contribute to gastric cancer development.(2, 5, 7, 8). The effect of PPI use on the gut microbiome may even be more prominent than the effects of antibiotics.(9) Among three recent meta-analyses on the topic, one found no association between long-term PPI-use and pre-malignant gastric lesions, based on six randomized controlled trials (1789 patients in total).(2) The second included an additional trial (2343 patients in total), and found no evidence of gastric tumor development in PPI-users with atrophy or intestinal metaplasia, while an increased risk of gastric hyperplasia was indicated.(6) The third, based on 11 observational studies (94,558 participants), reported a 40% increase of gastric cancer among PPI-users.(5) However, the impact of confounding by indication remains unknown. The present study aimed to assess the risk of gastric cancer in long-term PPI-users in a population-based design, while

1  
2  
3 taking confounding by indication for such treatment into account. For  
4  
5 comparison reasons, use of histamine 2 receptor antagonists (H2RAs), which  
6  
7 are used for similar indications as PPIs, was also studied.  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Methods**

***Design***

This was a nationwide Swedish population-based cohort study designed to examine the risk of gastric cancer in individuals exposed to maintenance therapy with PPIs (and to maintenance use of H2RAs), compared to the Swedish background population of the same sex, age and calendar period (7.1-7.6 million adults).(10)

Only adults (at least 18 years) without a history of any cancer were included. The participants were followed up from the first prescription of a PPI (or H2RA) during the period 1<sup>st</sup> July 2005 to 31<sup>st</sup> December 2012. The data were derived from high-quality and nationwide Swedish registries and information on individuals was linked between the registries by means of the unique Swedish personal identity number.(11) The source cohort included all Swedish residents who received at least one dispensed prescription of commonly prescribed drugs (listed in Appendix 1) between 1<sup>st</sup> July 2005 and 31<sup>st</sup> December 2014 (with follow-up for cancer until 31<sup>st</sup> December 2012). The study was approved by the Regional Ethical Review Board in Stockholm, and informed consent was not required (2014/1291-31/4).

***Patient involvement***

The Swedish patient organization for cancer of the esophagus, stomach, liver, and pancreas was involved in supporting the present study ([www.palema.org](http://www.palema.org)). The development of the research question and outcome measures were informed by patients' priorities, experience, and preferences. The results will be disseminated to study participants by means of patient

organizations. Patients are thanked in the acknowledgements.

### **Exposure**

The study exposure was maintenance therapy with a PPI (or an H2RA) according to the Swedish Prescribed Drug Registry, defined as a cumulative defined daily dose (DDD) of at least 6 months ( $\geq 180$  days) during the study period (before a potential cancer diagnosis). The DDD was the average maintenance dose per day for a drug used for its main indication in adults, which follows the World Health Organization (WHO) definition. This cumulative DDD was estimated by adding the DDD per package, which takes both the potency and the quantity of the drug into account, and is therefore a proxy for the duration of the exposure. The Anatomical Therapeutic Chemical classification system (ATC) was used to identify codes representing PPIs (code A02BC) and H2RAs (code A02BA). The participants were divided into 3 mutually exclusive exposure groups: (1) PPI-users ( $\geq 180$  days on PPIs and  $< 180$  days on H2RAs); (2) H2RA users ( $\geq 180$  days on H2RAs and  $< 180$  days on PPIs); and (3) PPI and H2RA users ( $\geq 180$  days on PPIs and  $\geq 180$  days on H2RAs). These medications were also available over-the-counter in Sweden, but only in small packages and at a much higher price per dose.<sup>(12)</sup> Thus, maintenance users had their medication prescribed and were therefore included in this study.

### **Outcome**

The main outcome was a first episode of any gastric cancer according to the Swedish Cancer Registry. Gastric cancer was defined by the diagnosis codes

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

C16.0 (cardia) and C16.1-C16.9 (non-cardia) in the 10<sup>th</sup> version of the International Classification of Diseases (ICD), and adenocarcinoma (dominating histological type) was defined by the histology code 096.

**Confounders**

Age, sex, and calendar period were adjusted for in the design. Confounding by indication was evaluated by analyzing indications for PPI-use separately (see Appendix 2 for ICD and ATC codes): gastro-esophageal reflux disease, Barrett’s esophagus, peptic ulcer disease, Zollinger-Ellison syndrome, gastro-duodenitis, dyspepsia/disruption of gastric function, *Helicobacter pylori* infection, *Helicobacter pylori* eradication (combined with *Helicobacter pylori* infection in the analyses), and maintenance use (≥180 days) of aspirin or other NSAIDs.

**Data sources**

Four nationwide Swedish registries, all maintained by the governmental National Board of Health and Welfare, provided data for the study.

**The Swedish Prescribed Drug Registry** started in 1<sup>st</sup> July 2005, and includes all drugs prescribed and dispensed in Sweden. Information from these prescriptions is transferred for registration monthly.(13) This registry is highly complete (patient identity data are missing in <0.3% of all items),(13) and was used to collect data on type of medication, dates of prescribing and dispensing, and DDD per package.

**The Swedish Cancer Registry** has at least a 96% complete registration of all cancers in Sweden since1958,(14) with 98% completeness for gastric

1  
2  
3 cancer.(15, 16) This registry was used to identify gastric cancer cases among  
4  
5 the cohort members and the corresponding background population (exact  
6  
7 annual numbers were available for the used age and sex categories), and to  
8  
9 exclude individuals with a history of any cancer.  
10

11 ***The Swedish Patient Registry*** has a full nationwide coverage for all  
12  
13 diagnoses from 1987.(11) The registration of out-patient specialist care is  
14  
15 nationwide since 2001. This registry was used to collect information on  
16  
17 indications for PPI-use.  
18  
19

20 ***The Swedish Causes of Death Registry*** has 100% completeness for  
21  
22 recording of dates of death since 1952.(17) This registry was used to collect  
23  
24 date of death.  
25  
26  
27  
28  
29

### 30 ***Statistical analyses***

31 All analyses were conducted according to a pre-planned protocol. The risk of  
32  
33 gastric cancer was compared between the exposed cohort and the entire  
34  
35 Swedish background population of the same sex (male or female), age group  
36  
37 (categorized as 18-39, 40-49, 50-59, 60-69 or  $\geq 70$  years), and calendar period  
38  
39 (2005-2006, 2007-2009 or 2010-2012). Standardized incidence ratios (SIRs)  
40  
41 and 95% confidence intervals (CIs) were calculated by dividing the observed  
42  
43 number of gastric cancer cases with the expected number, accounting for  
44  
45 changes in age and calendar categories.(18) The expected numbers were  
46  
47 derived from the Swedish Cancer Registry and Statistics Sweden.(10) Follow-  
48  
49 up time was calculated from the dispense date of the first prescription of PPI  
50  
51 (or H2RA) within the study period, until death, any cancer, or end of the study  
52  
53 period (December 31, 2012), whichever occurred first. Subgroup analyses  
54  
55  
56  
57  
58  
59  
60



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

were stratified for sex and age. Stratified analyses were performed for each indication whenever at least 10,000 exposed individuals were identified with this indication. To assess reverse causality (protopathic bias), a sensitivity analysis was conducted excluding all cancer cases occurring within one year of the start of the study. Duration of PPI-use was estimated based on the total DDD per package prescribed before any cancer diagnosis, and was categorized into <1.0 year, 1.0-2.9 years, 3.0-4.9 years or ≥5 years. The attributable fraction (AF) and the population attributable fraction (PAF) were calculated to estimate the proportion of gastric cancer in PPI-users and the total population, respectively (assuming causality). The following formulas were used:  $AF = (SIR - 1) / SIR$  and  $PAF = p(SIR - 1) / (p(SIR - 1) + 1)$ , where p represents the prevalence of PPI maintenance use in the population (based on the population size in 2009).<sup>(10)</sup> The Chi-square test was used to calculate p-values. There were no missing data on exposures, outcomes, age, sex or calendar period. When no information was available for the indication for PPI-use (25.0%), the indication was considered absent.

## Results

### *Study participants*

In total, 797,067 individuals on maintenance PPI therapy were included in the exposed cohort, resulting in 3,866,836 person-years of follow-up (mean 4.9 years). An additional 20,210 individuals had maintenance H2RA therapy, and 25,726 had both PPI and H2RA maintenance therapy. Figure 1 presents how the study participants were selected, and Table 1 shows their characteristics. Women constituted 58.5% of the PPI cohort, and a majority (66.1%) was younger than 70 years. Maintenance therapy with aspirin (34.8%) and NSAIDs (30.4%) were the most common indications for maintenance PPI use, followed by gastro-esophageal reflux (25.3%), gastro-duodenitis (13.2%) and peptic ulcer (10.0%). *Helicobacter pylori* was the indication in 7.3% of all participants, and dyspepsia in 5.5%. Barrett's esophagus and Zollinger-Ellison syndrome were rare indications (<1%) and therefore not eligible for separate analysis. A third (33.4%) of the participants had more than one of the listed indications, and was therefore included in more than one indication group. Overall mortality was slightly lower among PPI users (17.3%) compared to H2RA maintenance users (19.4%)( $p<0.001$ ), while mortality in gastric cancer patients did not differ significantly ( $p=0.782$ ).

### *Proton pump inhibitor use and overall risk of gastric cancer*

Among all participants exposed to maintenance PPIs, 2,219 (0.28%) developed gastric cancer during follow-up. Of these, 1,652 (74.4%) had non-cardia gastric cancer and 1,943 (87.6%) had adenocarcinoma (Table 1). Long-term PPI-users were at more than a 3-fold increased SIR of gastric

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

cancer of any type (SIR=3.38, 95% CI 3.25-3.53) and gastric adenocarcinoma (SIR=3.38, 95% CI 3.23-3.53) (Table 2). After excluding early gastric cancers, the risk remained increased (SIR=1.61, 95% CI 1.51-1.71). The risk of gastric cancer was highest among individuals receiving PPI for shorter than 1 year (SIR=12.82, 95% CI 12.19-13.47), but there was also evidence for an increased risk up to 3 years of PPI use, yet reduced risks for use over 5 years (Table 3). The strength of association was similar for cardia (SIR=3.55, 95% CI 3.27-3.86) and non-cardia gastric cancer (SIR=3.33, 95% CI 3.17-3.50). The risk of gastric cancer was similar in men (SIR=3.65, 95% CI 3.45-3.85) and women (SIR=3.07, 95% CI 2.87-3.28). The SIR was higher in younger ages; PPI-users younger than 40 years had SIR of 22.76 (95% CI 15.94-31.52) while those who were at least 70 years old had SIR of 2.76 (95% CI 2.61-2.92) (Table 2). The subgroup analyses for sex and age showed similar results as the overall findings in separate analyses for gastric adenocarcinoma, as well as for cardia and non-cardia gastric cancer (Table 2).

***Proton pump inhibitor use and risk of gastric cancer stratified by indication***

The SIR of gastric cancer was increased in each of the 10 studied groups of indications for PPI therapy (Table 4). The highest SIRs were found among participants exposed to indications with a known association with gastric cancer, i.e. *Helicobacter pylori* and peptic ulcer. However, the SIRs were also increased for indications without any such association. The SIR for gastric cancer was 3.04 (95% CI 2.80-3.31) for gastro-esophageal reflux. The SIR

were increased also for indications where a decreased risk of gastric cancer was expected. The SIR was 1.93 (95% CI 1.70-2.18) among those using PPIs because of aspirin use without any other indication. The SIRs were generally similar for non-cardia gastric cancer. Regarding cardia cancer, higher SIRs than for gastric cancer were found for the indication gastro-esophageal reflux and among those exposed to aspirin or other NSAIDs, while the associations for *Helicobacter pylori* and peptic ulcer were less strong (Table 4).

### ***Attributable risk***

Of all 5,823 gastric cancer patients diagnosed during the study period, 38.1% occurred among maintenance PPI-users. Based on the overall SIR of 3.38, and a prevalence of PPI-use of 10.7% among adults, the attributable fraction among PPI-users (AF) was 70.4%, and 20.3% in the total population (PAF), assuming causality. Using the lowest SIR according to Table 4 (SIR=1.41), the corresponding proportions were 29.1% (AF) and 4.2% (PAF), respectively. Among individuals younger than 40, the prevalence of PPI-use was 3.3%, and 37.1% of all 97 gastric cancers occurred among PPI-users, corresponding to AF of 95.6% and PAF of 41.8%.

### ***Use of histamine-2 receptor blockers only or proton pump inhibitors and histamine-2 receptor blockers and risk of gastric cancer***

There were 12 and 62 cases of gastric cancer among maintenance H2RAs users only and both H2RAs and PPIs, respectively. The risk of gastric cancer was not increased in the H2RA only group (SIR=0.57, 95% CI 0.29-0.99), and

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

was moderately increased in the group exposed to both PPIs and H2RAs  
(SIR=2.09, 95% CI 1.61-2.69).

For peer review only

## Discussion

This study provides some evidence of an increased risk of gastric cancer among maintenance PPI-users, including those who had PPI therapy for indications without any positive association with gastric cancer. Increased risks were found for both cardia and non-cardia gastric cancer, in both sexes and all age groups, although the risk was more pronounced in younger participants.

Strengths of the study include the nationwide and population-based design, the valid data on exposures, outcomes, and indications for PPI-use, and the large number of individuals exposed to maintenance treatment with PPIs. To our knowledge, this is the largest study to date on this topic,(19) and the cohort allowed analyses of differences in associations between underlying indications.

The main problems of this study are confounding, especially by indication, and reverse causality. Confounding by indication has been investigated by looking at the different indication groups, including groups without no increased risk of gastric cancer. Unfortunately, we could not identify any underlying indication for 25% of the PPI-users, and 39% of the H2RA-users. It is likely that clear indications were more readily recorded, while use for less rigorously diagnosed indications were more prone to be missing. This may indicate that PPI-users had more severe symptoms than H2RA-users, as also indicated by the higher proportion of PPI users presenting with multiple indications (34% vs. 19%). Yet, the lack of information on indication should not explain the associations for known indications – and those groups with no

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

apparent risk factors should be less likely to have severe gastrointestinal symptoms. Therefore, it was unexpected to still see such increased risks of gastric cancer in those groups. Unfortunately, we lacked information about some potential confounders, e.g. dietary factors, obesity, tobacco smoking and alcohol overconsumption since these are not collected in the nationwide Health Registries. However, the lack of association between H2RAs and gastric cancer argues against bias from confounding or selection from lifestyle factors or any other unknown factors. Since H2RAs were used for similar indications as PPIs (yet clearly became less popular), the lack of association between H2RAs and gastric cancer supports that the association between PPI-use and gastric cancer may be linked to the PPI medication *per se*. Yet, PPIs clearly became the first choice treatment for most indications for gastric acid suppression with almost 40 times more PPI maintenance users than H2RA maintenance users in Sweden. The high prevalence of PPI maintenance use among adults (10.7% in Sweden) also means that the large majority with recognized risk factors for gastric cancer will have received PPI treatment at some point. This clearly hampers assessment of PPI use as an independent risk factor and this may even be a larger problem in other study designs especially if indications of use cannot be assessed. The problem of reverse causality, i.e. individuals taking PPIs because of symptoms arising from an undetected cancer, should have been reduced by only including individuals with at least 180 days (6 months) of cumulative exposure before any cancer diagnosis. In addition, we excluded all individuals who had gastric cancer within a year after inclusion and also stratified the analyses by duration of use; which still showed increased risks (respectively

SIR=1.61 and SIR=2.19 among those with an exposure duration between 1 and 3 years). These increased risks may not be entirely explained by reverse causality and detection bias alone – since most cases of gastric cancer are believed to be detected within 1 year after onset of symptoms. There are no significant waiting times or socio-economic differences in access to endoscopy in Sweden, and no obvious differences were found between tumor stages (potential detection bias),(20) or anatomical locations between the PPI-exposed cohort and the background population (Appendix 3).

The seemingly decreasing risk estimates with a longer-duration of use is probably because PPI is beneficial for most individuals with known risk factors for cancer (e.g. peptic ulcers, *H. pylori*) – yet further research seems needed for those on maintenance therapy without gastrointestinal indications (e.g. aspirin and NSAIDs users).

Although our follow-up should ideally have been longer, our maximal follow-up time is 7.5 years, which is remarkably longer than all studies included in the 2 meta-analyses assessing pre-malignant lesions based on randomized controlled trials (6-36 months as maximal follow-up).(2, 6) Only 2 smaller cohort studies had longer follow-up,(1, 19) and these were included in the only meta-analysis evaluating gastric cancer risk based on observational studies.(5) We also limited our study to maintenance use, defined as at least 180 days of exposure, which is a commonly used approach.(6) Compliance and over-the-counter availability of some PPIs (Esomeprazole, Lansoprazole and Pantoprazole – only in pharmacy, not in retail sale) may have resulted in different actual dosages. H2RA were not available over-the-counter during the study period. Therefore, and because of lack of exposure data before the



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

study period, we cannot be certain that the registered dosage reflects the actual dosage.

The increased risks of gastric cancer among individuals with *Helicobacter pylori* and peptic ulcers were anticipated, since both these conditions are risk factors for gastric cancer, but the associations were stronger than expected, suggesting an additional etiological role of the PPI-use. Importantly, there was evidence of a substantially increased risk in individuals exposed to conditions not known to increase the risk of gastric cancer, including gastro-esophageal reflux. Moreover, increased risks were found even among participants without any gastrointestinal indications who were exposed only to maintenance treatment with aspirin or other NSAIDs, although these drugs are expected to decrease the risk of gastric cancer.(21, 22)

The higher risk of gastric cancer in the younger age group might be related to a recently described increase in atrophic gastritis in Sweden, as a potential consequence of the stabilizing sero-prevalence of *Helicobacter pylori* and increased prevalence of overweight and obesity.(23) Increasing risks of gastric cancer in young populations have been described previously, in particular of the diffuse type of gastric adenocarcinoma. (24, 25) Gastric symptoms in this group are also often over-looked, leading to a widespread (uncontrolled) use of PPIs.(25) Additionally, gastric carcinogenesis may be accelerated in younger patients (who also have a higher likelihood of a family history of gastric cancer),(26) which may in turn contribute to a higher vulnerability to PPI's potentially harmful effects. A further exploration into the

presence and severity of gastrointestinal risk factors, age-specific effects and potential mechanistic pathways is warranted.

The difference in the association with gastric cancer following maintenance therapy with PPIs and H2RAs could be explained simply by the fact that PPIs are more potent than H2RAs in inhibiting acid secretion.(27) PPIs block the gastric proton pumps, while H2RAs compete with histamine for the histamine-2 receptors. At recommended dosages, PPIs induce a more profound and prolonged acid inhibition than H2RAs. Moreover, the acid inhibition of PPIs also increases over time, while the effects of H2RAs fade.(27)

There are several mechanisms that might explain an association between PPI-use and gastric cancer. Already 30 years ago, animal studies showed that profound inhibition of gastric acid secretion in rodents induces gastric tumours, with secondary over-stimulation (hypergastrinemia) leading to enterochromaffin-like cell (ECL) hyperplasia as generally accepted mechanism of this carcinogenic effect.(28, 29) ECL differentiation has also been described in human gastric carcinomas, particularly in the signet ring subtype.(30, 31) Histopathological changes occur due to the blockage of the normal gastric acid secretion, leading to hypergastrinemia that might cause hyperproliferation of the gastric mucosa, chronic hypochlorhydria (reduction of hydrochloric acid in the gastric juice), chronic inflammation and disappearance of normal mucosal glands and their replacement by intestinal glands, and possible gastric atrophy.(6) Long-term PPI use has also been linked to the development of fundic gland polyps.(32) All these mucosal

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

changes could promote gastric carcinogenesis. The blocked gastric acid secretion could decrease physiological defense mechanisms against pathogenic bacteria.(7, 8) Diarrhea is for example a well-known side-effect of PPI-use, which is often due to infection with the bacteria *Clostridium difficile*, non-thyphoid *Salmonella species* or *Campylobacter jejuni*.(7) This reduced bacterial defense mechanism might, in the long term, result in chronic inflammation and eventually gastric cancer development. Decreased gastric acidity may also result in increased bacterial colonization, including non-gastric microorganisms, and a greater number of bacteria that produce nitrosamines, which are well-established carcinogens of the gastric mucosa.(5, 33) The impact of these proposed underlying mechanisms may also be person- and time-dependent, which could explain why PPIs are seemingly more harmful in for example the younger age groups.

The results of a single study, even of this size, cannot determine causality. Yet, the effect size, consistency across indications (especially those not associated with an increased risk of gastric cancer), lack of association with H2RAs, and plausible carcinogenic mechanisms mean that maintenance PPI-use cannot be dismissed as a potentially independent risk factor for gastric cancer. A higher awareness for gastric cancer could potentially be considered in maintenance users of PPI, especially if known underlying risk factors are present. Thus, these findings need confirmation in future investigations, especially considering the generalizability of the results to populations with higher incidence of gastric cancer or other distribution of risk factors.

To conclude, this large and population-based cohort study provides evidence of a substantially increased risk of gastric cancer following maintenance use of PPIs *per se*. The consistency across sexes, age groups, and indication groups, including indications that do not increase the risk of gastric cancer, support the overall finding. If confirmed in further research, especially among those without gastrointestinal risk factors, these findings challenge a broad use of long-term PPI-use, particularly for conditions where the indication is weak.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Acknowledgements:** We are grateful to all thousands of patients, physicians and healthcare staff members who contributed to the data collection, and to the Swedish National Board of Health and Welfare that collected the data. NB had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. NB conducted and is responsible for the data-analysis. Everyone who contributed significantly to the work is listed as co-author.

**Funding source:** The work was supported by the Swedish Research Council (839-2008-7496), Swedish Cancer Society (CAN 2015/460), the Karolinska Institutet Distinguished Professor Award to Jesper Lagergren (D-02418/2010) and SFO Epidemiology (Young Scholar Grant to Nele Brusselaers).

**Competing interest statement:** There are no competing interests.

**Data sharing statement:** We are willing to share data upon request after ethical approval has been approved by the relevant committee and the governmental agencies that maintain the data.

**References**

1. Moller H, Nissen A, Mosbech J. Use of cimetidine and other peptic ulcer drugs in Denmark 1977-1990 with analysis of the risk of gastric cancer among cimetidine users. Gut. 1992 Sep;33(9):1166-9. PubMed PMID: 1358764. Pubmed Central PMCID: PMC1379479.

2. Eslami L, Nasser-Moghaddam S. Meta-analyses: does long-term PPI use increase the risk of gastric premalignant lesions? Arch Iran Med. 2013 Aug;16(8):449-58. PubMed PMID: 23906249. Epub 2013/08/03. eng.
3. Shin JM, Sachs G. Pharmacology of proton pump inhibitors. Curr Gastroenterol Rep. 2008 Dec;10(6):528-34. PubMed PMID: 19006606. Pubmed Central PMCID: PMC2855237.
4. The World Medicines Situation (WMS) 2011 - Pharmaceutical consumption - Annex 4 Top 5 medicine classes by volume and country income category.
5. Ahn JS, Eom CS, Jeon CY, Park SM. Acid suppressive drugs and gastric cancer: a meta-analysis of observational studies. World J Gastroenterol. 2013 Apr 28;19(16):2560-8. PubMed PMID: 23674860. Pubmed Central PMCID: 3646149. Epub 2013/05/16. eng.
6. Song H, Zhu J, Lu D. Long-term proton pump inhibitor (PPI) use and the development of gastric pre-malignant lesions. Cochrane Database Syst Rev. 2014;12:CD010623. PubMed PMID: 25464111.
7. Canani RB, Terrin G. Gastric acidity inhibitors and the risk of intestinal infections. Curr Opin Gastroenterol. 2010 Jan;26(1):31-5. PubMed PMID: 19907324.
8. Seto CT, Jeraldo P, Orenstein R, Chia N, DiBaise JK. Prolonged use of a proton pump inhibitor reduces microbial diversity: implications for Clostridium difficile susceptibility. Microbiome. 2014;2:42. PubMed PMID: 25426290. Pubmed Central PMCID: PMC4242847.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

9. Imhann F, Bonder MJ, Vich Vila A, Fu J, Mujagic Z, Vork L, et al. Proton pump inhibitors affect the gut microbiome. Gut. 2016 May;65(5):740-8. PubMed PMID: 26657899. Pubmed Central PMCID: 4853569.

10. Statistics Sweden - Statistical Database Population Statistics (accessed 06 April 2016) <http://www.statistikdatabasen.scb.se/>. Available from: <http://www.statistikdatabasen.scb.se/>.

11. Ludvigsson JF, Andersson E, Ekbom A, Feychting M, Kim JL, Reuterwall C, et al. External review and validation of the Swedish national inpatient register. BMC Public Health. 2011;11:450. PubMed PMID: 21658213. Pubmed Central PMCID: 3142234. Epub 2011/06/11. eng.

12. FASS (Farmaceutiska specialiteter i Sverige/ Pharmaceutical specialities in Sweden) [Internet]. 2015 [cited November 2016]. Available from: [www.fass.se](http://www.fass.se).

13. Wettermark B, Hammar N, Fored CM, Leimanis A, Otterblad Olausson P, Bergman U, et al. The new Swedish Prescribed Drug Register--opportunities for pharmacoepidemiological research and experience from the first six months. Pharmacoepidemiol Drug Saf. 2007 Jul;16(7):726-35. PubMed PMID: 16897791. Epub 2006/08/10. eng.

14. Barlow L, Westergren K, Holmberg L, Talback M. The completeness of the Swedish Cancer Register: a sample survey for year 1998. Acta Oncol. 2009;48(1):27-33. PubMed PMID: 18767000. Epub 2008/09/04. eng.

15. Lindblad M, Ye W, Lindgren A, Lagergren J. Disparities in the classification of esophageal and cardia adenocarcinomas and their influence on reported incidence rates. Ann Surg. 2006 Apr;243(4):479-85. PubMed PMID: 16552198. Pubmed Central PMCID: 1448962. Epub 2006/03/23. eng.

16. Ekstrom AM, Signorello LB, Hansson LE, Bergstrom R, Lindgren A, Nyren O. Evaluating gastric cancer misclassification: a potential explanation for the rise in cardia cancer incidence. *J Natl Cancer Inst.* 1999 May 5;91(9):786-90. PubMed PMID: 10328109. Epub 1999/05/18. eng.
17. The National Board of Health and Welfare, Sweden. Causes of death in Sweden 2009. 2011. Available from: <http://www.socialstyrelsen.se/publikationer2011/2011-3-22>.
18. Breslow N, Day N. *Statistical Methods in Cancer Research: The design and analysis of cohort studies.* Lyon1987.
19. Poulsen AH, Christensen S, McLaughlin JK, Thomsen RW, Sorensen HT, Olsen JH, et al. Proton pump inhibitors and risk of gastric cancer: a population-based cohort study. *Br J Cancer.* 2009 May 5;100(9):1503-7. PubMed PMID: 19352380. Pubmed Central PMCID: PMC2694435.
20. Sobin L, Wittekind C. *International Union Against Cancer (UICC)- 6th edition: Classification of malignant tumours.* 6th ed2002.
21. Gurpinar E, Grizzle WE, Piazza GA. NSAIDs inhibit tumorigenesis, but how? *Clin Cancer Res.* 2014 Mar 1;20(5):1104-13. PubMed PMID: 24311630. Pubmed Central PMCID: PMC3947450.
22. Ford AC. Chemoprevention for gastric cancer. *Best Pract Res Clin Gastroenterol.* 2011 Aug;25(4-5):581-92. PubMed PMID: 22122773. Epub 2011/11/30. eng.
23. Song H, Held M, Sandin S, Rautelin H, Eliasson M, Soderberg S, et al. Increase in the Prevalence of Atrophic Gastritis Among Adults Age 35 to 44 Years Old in Northern Sweden Between 1990 and 2009. *Clin Gastroenterol Hepatol.* 2015 Sep;13(9):1592-600 e1. PubMed PMID: 25857683.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

24. Anderson WF, Camargo MC, Fraumeni JF, Jr., Correa P, Rosenberg PS, Rabkin CS. Age-specific trends in incidence of noncardia gastric cancer in US adults. *JAMA*. 2010 May 05;303(17):1723-8. PubMed PMID: 20442388. Pubmed Central PMCID: 3142962.

25. Pisanu A, Podda M, Cois A, Uccheddu A. Gastric cancer in the young: is it a different clinical entity? A retrospective cohort study. *Gastroenterology research and practice*. 2014;2014:125038. PubMed PMID: 24688534. Pubmed Central PMCID: 3943291.

26. Tavares A, Gandra A, Viveiros F, Cidade C, Maciel J. Analysis of clinicopathologic characteristics and prognosis of gastric cancer in young and older patients. *Pathol Oncol Res*. 2013 Jan;19(1):111-7. PubMed PMID: 22573278.

27. Lamers CB. The changing role of H2-receptor antagonists in acid-related diseases. *Eur J Gastroenterol Hepatol*. 1996 Oct;8 Suppl 1:S3-7. PubMed PMID: 8930573.

28. Havu N. Enterochromaffin-like cell carcinoids of gastric mucosa in rats after life-long inhibition of gastric secretion. *Digestion*. 1986;35 Suppl 1:42-55. PubMed PMID: 3792671.

29. Poynter D, Pick CR, Harcourt RA, Selway SA, Ainge G, Harman IW, et al. Association of long lasting unsurmountable histamine H2 blockade and gastric carcinoid tumours in the rat. *Gut*. 1985 Dec;26(12):1284-95. PubMed PMID: 2867954. Pubmed Central PMCID: 1433126.

30. Bakkelund K, Fossmark R, Nordrum I, Waldum H. Signet ring cells in gastric carcinomas are derived from neuroendocrine cells. *The journal of*

1  
2  
3 histochemistry and cytochemistry : official journal of the Histochemistry  
4  
5 Society. 2006 Jun;54(6):615-21. PubMed PMID: 16344325.  
6

7 31. Waldum HL, Aase S, Kvetnoi I, Brenna E, Sandvik AK, Syversen U, et  
8  
9 al. Neuroendocrine differentiation in human gastric carcinoma. Cancer. 1998  
10  
11 Aug 01;83(3):435-44. PubMed PMID: 9690535.  
12  
13

14 32. Ally MR, Veerappan GR, Maydonovitch CL, Duncan TJ, Perry JL,  
15  
16 Osgard EM, et al. Chronic proton pump inhibitor therapy associated with  
17  
18 increased development of fundic gland polyps. Dig Dis Sci. 2009  
19  
20 Dec;54(12):2617-22. PubMed PMID: 19830554.  
21  
22

23 33. Engstrand L, Lindberg M. Helicobacter pylori and the gastric  
24  
25 microbiota. Best Pract Res Clin Gastroenterol. 2013 Feb;27(1):39-45.  
26  
27 PubMed PMID: 23768551. Epub 2013/06/19. eng.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Figure 1: Flowchart of the construction of the cohort exposed to maintenance therapy with proton pump inhibitors (PPIs) and/or H2 receptor antagonists (H2RAs).**

For peer review only

**Table 1: Descriptive characteristics of the study cohort exposed to maintenance therapy with proton pump inhibitors (PPIs) and/or H2 receptor antagonists (H2RAs).**

		PPIs only Number (%)	H2RAs only Number (%)	PPIs and H2RAs Number (%)
<b>Total</b>		797,067	20,210	25,726
<b>Sex</b>				
	<i>Men</i>	330,885 (41.5)	8,076 (40.0)	9,778 (38.0)
	<i>Women</i>	466,182 (58.5)	12,134 (60.0)	15,948 (62.0)
<b>Age (years)</b>				
	<40	89,231 (11.2)	1,885 (9.3)	1,822 (7.1)
	40-49	104,003 (13.1)	2,461 (12.2)	3,268 (12.7)
	50-59	155,963 (19.6)	3,971 (19.7)	5,878 (22.9)
	60-69	177,606 (22.3)	4,563 (22.6)	6,354 (24.7)
	≥70	270,264 (33.9)	7,330 (36.3)	8,404 (32.7)
<b>Calendar period</b>				
	2005-2006	446,068 (56.0)	16,550 (81.9)	23,066 (90.0)
	2007-2009	222,257 (27.9)	2,670 (13.2)	2,304 (9.0)
	2010-2012	128,742 (16.2)	990 (4.9)	356 (1.4)
<b>Indications</b>				
	<i>Gastro-esophageal reflux</i>	201,868 (25.3)	2,073 (10.3)	6,727 (26.2)
	<i>Barrett's esophagus</i>	6,044 (0.8)	17 (0.1)	172 (0.7)
	<i>Peptic ulcers</i>	79,597 (10.0)	1,226 (6.1)	2,633 (10.2)
	<i>Zollinger-Ellison syndrome</i>	31 (0.0)	0 (0.0)	3 (0.01)
	<i>Gastro-duodenitis</i>	104,955 (13.2)	1,321 (6.5)	3,695 (14.4)
	<i>Dyspepsia</i>	43,901 (5.5)	649 (3.2)	1,664 (6.5)
	<i>Helicobacter pylori</i>	58,366 (7.3)	641 (3.2)	2,041 (7.9)
	<i>Long-term NSAID use</i>	241,958 (30.4)	5,012 (24.8)	9,628 (37.4)
	<i>Long-term aspirin use</i>	277,128 (34.8)	6,533 (32.3)	9,929 (38.6)
<b>Number of indications</b>				
	0	199,608 (25.0)	7,919 (39.2)	5,402 (21.0)
	1	330,027 (41.4)	8,487 (42.0)	10,320 (40.1)
	≥2	267,432 (33.6)	3,804 (18.8)	10,004 (38.9)
<b>Gastric cancer</b>				
	<i>All</i>	2,219 (0.28)	12 (0.06)	62 (0.24)
	<i>Adenocarcinoma</i>	1,943 (0.24)	10 (0.05)	54 (0.21)
	<i>Cardia</i>	567 (0.07)	1 (0.00)	13 (0.05)
	<i>Non-cardia</i>	1,652 (0.21)	11 (0.05)	49 (0.19)
<b>Years of follow-up</b>				
	Total	3,866,836	116,015	163,519
	Mean	4.9	5.7	6.4
<b>Cumulative defined daily dosages (DDD), estimated number of days</b>				
	Median	856	450	1741
	Interquartile range	356-2102	270-990	1043-2732
<b>Mortality</b>				
	Total	137,619 (17.3)	3,915 (19.4)	3,812 (14.8)
	In gastric cancer cases	1,473 (66.4)	9 (75.0)	39 (62.9)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

**Table 2: Standardized incidence ratios (SIRs) by age, sex and calendar period and 95% confidence intervals (CIs) of gastric cancer in all individuals exposed to proton pump inhibitors (PPIs), stratified by sex and age-group.**

	Gastric cancer		Gastric adenocarcinoma		Cardia cancer		Non-cardia gastric cancer	
	Number of cases	SIR (95% CI)	Number of cases	SIR (95% CI)	Number of cases	SIR (95% CI)	Number of cases	SIR (95% CI)
Total	2,219	3.38 (3.25-3.53)	1,943	3.38 (3.23-3.53)	567	3.55 (3.27-3.86)	1,652	3.33 (3.17-3.50)
Sex								
Men	1,296	3.65 (3.45-3.85)	1,166	3.66 (3.45-3.88)	437	3.84 (3.48-4.21)	859	3.56 (3.33-3.81)
Women	923	3.07 (2.87-3.28)	777	3.02 (2.82-3.25)	130	2.84 (2.38-3.38)	793	3.11 (2.99-3.34)
Age, years								
<40	36	22.76 (15.94-31.52)	27	25.30 (16.67-36.80)	7	27.33 (10.95-56.30)	29	21.88 (14.65-31.43)
40-49	94	7.36 (5.95-9.01)	76	7.76 (6.11-9.71)	24	7.54 (4.83-11.22)	70	7.30 (5.69-9.23)
50-59	303	5.63 (5.01-6.30)	263	5.82 (5.14-6.57)	94	5.50 (4.44-6.72)	209	5.69 (4.94-6.51)
60-69	576	3.85 (3.54-4.18)	501	3.81 (3.49-4.16)	191	3.94 (3.40-4.54)	385	3.81 (3.44-4.21)
≥70	1,210	2.76 (2.61-2.92)	1,076	2.77 (2.61-2.94)	251	2.77 (2.44-3.14)	959	2.76 (2.59-2.94)

**Table 3: Standardized incidence ratios (SIRs) by estimated duration of use and 95% confidence intervals (CIs) of gastric cancer in all individuals exposed to proton pump inhibitors (PPIs)**

Duration	Number of cases /	
	Total	SIR (95% CI)
< 1.0 year	1,552	12.82 (12.19-13.47)
1.0-2.9 years	2,193	2.19 (1.98-2.42)
3.0-4.9 years	1,098	1.10 (0.91-1.31)
≥ 5.0 years	153	0.61 (0.52-0.72)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

**Table 4: Standardized incidence ratios (SIRs) by age, sex and calendar period and 95% confidence intervals (CIs) of gastric cancer in all individuals exposed to proton pump inhibitors (PPIs), stratified by indication.**

	Gastric cancer		Cardia cancer		Non-cardia gastric cancer	
	Number of cases / total	SIR (95% CI)	Number of cases / total	SIR (95% CI)	Number of cases / total	SIR (95% CI)
All	2,219/ 797,067	3.38 (3.25-3.53)	567/ 797,067	3.55 (3.27-3.86)	1652/ 797,067	3.33 (3.17-3.50)
Gastro-esophageal reflux	557/ 201,868	3.04 (2.80-3.31)	175/ 201,868	3.84 (3.29-4.45)	382/ 201,868	2.78 (2.51-3.07)
Peptic ulcers	721/ 79,597	8.75 (8.12-9.41)	71/ 79,597	3.51 (2.74-4.43)	650/ 79,597	10.45 (9.66-11.28)
Gastro-duodenitis	350/ 104,955	3.68 (3.31-4.09)	63/ 104,955	2.71 (2.09-3.47)	287/ 104,955	3.99 (3.55-4.48)
Dyspepsia	136/ 43,901	3.07 (2.58-3.63)	33/ 43,901	3.10 (2.13-4.35)	103/ 43,901	3.07 (2.50-3.72)
<i>Helicobacter pylori</i>	440/ 58,366	9.76 (8.87-10.71)	48/ 58,366	4.19 (3.09-5.55)	392/ 58,366	11.67 (10.53-12.87)
No <i>Helicobacter pylori</i>	1,779/ 781,955	2.91 (2.78-3.05)	519/ 781,955	3.50 (3.21-3.82)	1,302/ 781,955	2.73 (2.58-2.88)
NSAIDs exposure	347/ 256,598	1.82 (1.64-2.03)	94/ 256,598	2.07 (1.67-2.53)	253/ 256,598	1.75 (1.54-1.97)
Only NSAIDs exposure	93/ 118,987	1.41 (1.14-1.73)	36/ 118,987	2.29 (1.60-3.17)	57/ 118,987	1.14 (0.86-1.48)
Aspirin exposure	872/ 293,590	2.42 (2.26-2.60)	211/ 293,590	2.73 (2.37-3.13)	588/ 293,590	2.33 (2.14-2.53)
Only aspirin exposure	252/ 126,883	1.93 (1.70-2.18)	85/ 126,883	2.73 (2.18-3.37)	169/ 126,883	1.68 (1.43-1.95)

\* NSAIDs, non-steroidal anti-inflammatory drugs. One individual could have different indications, except for the group exposed only to NSAIDs or aspirin.




Figure 1: Flowchart of the construction of the cohort exposed to maintenance therapy with proton pump inhibitors (PPIs) and/or H2 receptor antagonists (H2RAs).

190x209mm (300 x 300 DPI)



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Appendix 1: Description of original cohort, the “Chemoprevention of Cancer” cohort.**

This cohort included all individuals residing in Sweden who received at least one dispensed prescription of one of the following commonly prescribed drugs between July 2005 and December 2014 (with corresponding ATC codes): sex hormones (G03), drugs for peptic ulcers and gastro-esophageal reflux disease (A02B), acetylsalicylic acid (B01AC06, N02BA), non-steroidal anti-inflammatory drugs (M01A), HMG CoA reductase inhibitors (C10AA, C10BA), drugs affecting bone structure and mineralization, (M05B), and antibiotics (J01AA, J01CA04, J01FA, J01MA, J01XD, J01XE, J04AB04). This cohort included approximately 85% of all Swedish residents, with especially high coverage of adults. All adult individuals receiving at least one dispensed prescription of a gastric acid inhibitor (proton pump inhibitor or H2 receptor antagonist) were included.

**Appendix 2: International Classification of Diseases (ICD) and Anatomical Therapeutic Chemical classification system (ATC) codes to describe the indication for gastric acid inhibitors.**

	ICD-7 (Since 1955)	ICD-8 (Since 1968)	ICD-9 (Since 1987)	ICD-10 (Since 1997)
Reflux codes	784,3; 539,11- 539,12; 560,4	530,93- 530,94; 551,3; 784,3	530B-C; 553D; 787B	K20-21; K44; R12
Barrett's esophagus	-	-	-	K227
Peptic ulcer disease	540-541	531-533	531-533	K25-K27
Zollinger-Ellison syndrome / Hypergastrinemia	-	-	251F	E16.4
Gastro-duodenitis	543	535	535	K29
Dyspepsia/disruption of gastric function	544	536	536	K30-K31
<i>Helicobacter pylori</i> infection	-	-	-	B96.8 or B98.0
<b>ATC</b>				
Non-steroidal anti-inflammatory drugs	M01A			
Aspirin	B01AC06, N02BA			
<i>Helicobacter pylori</i> eradication		A02BD, or A02BC (PPI) in combination with 2 out of 3 antibiotics (Amoxicillin [J01CA04]; Clarithromycin [J01FA09], or metronidazole [J01XD]); or combined with metronidazole [J01XD], doxycycline [J01AA02] and bismuth [A02BX05]; or with amoxicillin [J01CA04] and levofloxacin [J01MA12].		

Note: an indication was considered present if the ICD code occurred at least once in the in- or out-Patient Registry. NSAIDs and Aspirin were only considered if the cumulative dosage was  $\geq 180$  days.

**Appendix 3: Tumor stage and anatomical locations of gastric cancer in the cohort exposed to maintenance use of proton pump inhibitors (PPIs) and in the Swedish background population of the same age.**

	PPI maintenance cohort	Background population
	Number (%)	Number (%)
<b>Total</b>	2219 (100.0)	5821 (100.0)
<b>Tumor stage - Cardia</b>		
0-1	72 (12.7)	134 (8.9)
2	36 (6.4)	80 (5.3)
3	207 (36.5)	480 (31.9)
4	145 (25.6)	496 (33.0)
unclear/incomplete	107 (18.9)	315 (20.9)
<b>Tumor stage - Non-cardia</b>		
0-1	291 (17.6)	617 (14.3)
2	68 (4.1)	168 (3.9)
3	408 (24.7)	979 (22.7)
4	422 (25.5)	1260 (29.2)
unclear/incomplete	463 (28.0)	1292 (29.9)
<b>Anatomical location</b>		
Cardia	567 (25.6)	1505 (25.9)
Fundus	97 (4.4)	268 (4.6)
Corpus	389 (17.5)	1040 (17.9)
Antrum	398 (17.9)	915 (15.7)
Pylorus	157 (7.1)	303 (5.2)
Minor curvature	74 (3.3)	173 (3.0)
Major curvature	33 (1.5)	93 (1.6)
Overgrowth of unknown origin	2 (0.1)	15 (0.3)
Unspecified	502 (22.6)	1509 (25.9)

STROBE Statement—Checklist of items that should be included in reports of *cohort studies*

	Item No	Recommendation	Page
<b>Title and abstract</b>	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	p 1-2 (abstract & title)
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	p 2
<b>Introduction</b>			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	p 4-5
Objectives	3	State specific objectives, including any prespecified hypotheses	p 4-5
<b>Methods</b>			
Study design	4	Present key elements of study design early in the paper	p 6-10
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	p 6-10
Participants	6	(a) Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	p 6-10
		(b) For matched studies, give matching criteria and number of exposed and unexposed	n/a
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	p 6-10 + appendix
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	p 6-10
Bias	9	Describe any efforts to address potential sources of bias	p 6-10
Study size	10	Explain how the study size was arrived at	Fig 1
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	p 6-10
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	p 9-10
		(b) Describe any methods used to examine subgroups and interactions	p 9-10
		(c) Explain how missing data were addressed	p 9-10
		(d) If applicable, explain how loss to follow-up was addressed	n/a
		(e) Describe any sensitivity analyses	p 9-10
<b>Results</b>			
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	Fig 1
		(b) Give reasons for non-participation at each stage	Fig 1
		(c) Consider use of a flow diagram	Fig 1
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	p 11, Table 1
		(b) Indicate number of participants with missing data for each variable of interest	n/a
		(c) Summarise follow-up time (eg, average and total amount)	Table 1
Outcome data	15*	Report numbers of outcome events or summary measures over time	Tables 1-3
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted	Tables 2-3

		estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	
		(b) Report category boundaries when continuous variables were categorized	Table 1-2
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	Discussion p 13
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	p 11-14
<b>Discussion</b>			
Key results	18	Summarise key results with reference to study objectives	p 15-19
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	p 15-19
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	p 15-19
Generalisability	21	Discuss the generalisability (external validity) of the study results	p 15-19
<b>Other information</b>			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	p 3, p 10

\*Give information separately for exposed and unexposed groups.

**Note:** An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at <http://www.strobe-statement.org>.